Vol. 4. No. 3. September, 1947.

THE QUEENSLAND JOURNAL OF AGRICULTURAL SCIENCE

COMMONWEALTH INST
ENTOMOLOGY LIBRARY
-7OCT
SERIAL Aus.40
SEPARATE

CONTENTS

PAGE.

Studies in the Environment of Queensland. 1, The Climatology of Semi-Arid Pastoral Areas. By Joan N. Farmer, S. L. Everist and G. R. Moule 21

Electrocardiograms of Young Lambs. By W. T. Kelso 60

PUBLISHED QUARTERLY

By the Queensland Department of Agriculture and Stock, Brisbane

Issued by Direction of Hon. H. H. COLLINS,
Secretary for Agriculture and Stock.

EDITORIAL BOARD.

Editor: C. W. WINDERS, B.Sc.Agr.

J. LEGG, B.Sc., D.V.Sc., M.R.C.V.S., Director of Research, Division of Animal Industry;

C. J. McKEON, Q.D.A., Director of Agriculture, Agriculture Branch, Division of Plant Industry;

J. F. F. REID, Editor of Departmental Publications;

J. H. SIMMONDS, M.B.E., M.Sc., Officer-in-Charge and Senior Pathologist, Science Branch, Division of Plant Industry;

J. H. SMITH, M.Sc., N.D.A., Senior Entomologist, Science Branch, Division of Plant Industry;

W. A. T. SUMMERVILLE, D.Sc., Director, Division of Plant Industry;

R. VEITCH, B.Sc.Agr., B.Sc.For., F.R.E.S., Assistant Under Secretary (Technical);

M. WHITE, M.Sc., Ph.D., A.A.C.I., Agricultural Chemist and Bio-Chemist, Chemical Laboratory, Division of Plant Industry.

The Journal is supplied free, on request, to scientific agricultural and veterinary institutions and organizations. Application should be made to the Under Secretary, Department of Agriculture and Stock, Brisbane, Queensland.

THE QUEENSLAND JOURNAL OF AGRICULTURAL SCIENCE

VOL. 4. NO. 3.

SEPTEMBER, 1947

STUDIES IN THE ENVIRONMENT OF QUEENSLAND. 1, THE CLIMATOLOGY OF SEMI-ARID PASTORAL AREAS.

By JOAN N. FARMER, B.Sc., Technical Assistant, Sheep and Wool Branch; S. L. EVERIST, B.Sc., Botanist, Science Branch; and G. R. MOULE, B.V.Sc., Officer-in-Charge, Sheep and Wool Branch.

SUMMARY.

The literature dealing with climatic classification is reviewed, and the P/E ratio used as a method of measuring effective rainfall.

The seasonal incidence of effective rainfall is calculated for bi-monthly periods, and zonation is undertaken on the basis of its reliability.

Summer and winter P/E differences, mean minimum winter and mean maximum summer temperatures are considered and the broad influence of climatic factors on the growth and distribution of plants and on sheep husbandry is discussed.

INTRODUCTION.

The study of climate is of paramount importance in a primary producing country where the production of wool, meat and milk depends largely upon natural vegetation. Different climatic zones produce vegetation specific to themselves and thus the type of primary production best suited to any given region depends largely upon its climate.

In a pastoral zone methods of animal husbandry often make the difference between success and failure. This is especially true in the raising of Merino sheep. Wool is expensive to produce and it is important that the utmost efficiency be attained in its production. To do this it is essential that the influence of environment be thoroughly understood. It is essential, also, to know how far results obtained from different methods of husbandry in one district can be applied to others, and to determine the limits of economic wool production and economic sheep-breeding.

The results of a study which has been made of the climatic data from the semi-arid region of Queensland are presented in this paper. It is obvious that considerable differences exist in the climate of various parts of the main

sheep grazing areas of the State, and it is thought that these differences may be correlated with natural vegetation, which in turn governs the growth and reproduction of sheep as well as wool production.

LITERATURE.

Various criteria have been used for the classification of climate and the different methods have been reviewed by Moreau (1938) and Prescott (1934), who indicate the limitations and applications of the classifications offered by a number of writers, including those of Köppen, Thornthwaite, de Martonne, Emberger, Meyer, Lang, Prescott, Davidson and Trumble.

Köppen (see Moreau, 1938, and Lawrence, 1937) divided the world into climatic zones using temperature, rainfall and rainfall distribution. Thornthwaite (1931, 1933; see also Moreau, 1938, and Prescott, 1934) obtained his "precipitation efficiency index" as a complicated function of rainfall and temperature, but took no account of relative humidity. Both of these methods are useful over large areas, but are not suitable for application to a particular region, such as the pastoral areas of Queensland.

Andrews and Maze (1933) used a modification of de Martonne's "Index of Aridity" ($\frac{P}{T + 10}$, where P = precipitation in cms., T = °C.) in Australia with some success. Their method was to classify climates on the basis of the number of dry months in the year, the criterion of a dry month being an index of less than 1. Again, humidity was not taken into account. It seems that, where wet and dry bulb readings are available, the humidity should be considered, its controlling effect on evaporation being obvious.

Davidson (1933, 1935) and Prescott (1934, 1936) investigated the Meyer Ratio $\frac{P}{S.D.}$ (where P = precipitation; S.D. = saturation deficit, which is the difference between the saturation vapour pressure of water at a given temperature and the partial pressure of water vapour actually present in the environment at that temperature). Davidson (1934, 1935) applied the supposition that evaporation from a free water surface is proportional to saturation deficit, provided all other factors are constant. Actually wind velocity, atmospheric pressure, insolation, turbulence, exposure, and nature of land surface have an effect on evaporation. Prescott's investigation of 144 monthly values at a selected range of 12 Australian stations (1938) showed a correlation coefficient of ·972 between evaporation and saturation deficit. Davidson (1933) showed that for South Australia mean monthly values of S.D. could be expressed in terms of evaporation by referring to data for a free water surface at Adelaide and he formulated the equation $\frac{P}{E} = m \frac{P}{S.D.}$ (where m is a constant). Prescott (1938) worked with this conversion factor, using the formula E = 21.2 S.D. for a month of 30 days. The same numerical relationship was used by Lawrence (1941) in an analysis of the climate of tropical Australia.

Prescott (1934; see also Moreau, 1938) noted the weakness of using maximum and minimum temperatures for the calculation of saturation deficit and stated that the minimum is less important than the dew point. In semi-arid

regions this is true to some extent but it is not as important as in more humid areas, since on a very large number of days during the year minimum temperatures do not fall below the dew point. This applies particularly to Queensland, where the air masses in winter are mostly of the dry continental type.

Prescott (1942, 1943) also pointed out the existence of a difference in phase between temperature and evaporation. This difference shows a maximum of 21 days at the ocean coast and falls off rapidly to values of the order of seven days within 150 miles of the coast. The only data available are on a monthly basis so it was not found practicable by the authors to apply any correction for phase lag, and it was assumed that monthly evaporation is a function of the temperature and humidity recorded for the same month.

Prescott (1938) concluded that, to find some expression for the climatic conditions of a given environment, temperature, rainfall and humidity must be linked with soil moisture conditions, and the P/E Transeau ratio appeared to be a practical measure of effective soil moisture, E being calculated from temperature and humidity. Trumble (1939) considered a value of P/E = ·3 (computed from total rainfall and evaporation for any one month) as being effective for producing plant growth in South Australia. The difference in values required for South Australia and tropical Australia were noted by Lawrence (1941). In her study of climate in the latter part of the continent she used P/E = ·25 as the minimum ratio limiting a month of effective rainfall.

Szymkiewiez (1925) suggested the substitution of $\text{S.D.} \times \frac{273 + T}{273}$ for S.D. in Prescott's equation, given above, thus taking into consideration the greater diffusivity of water vapour at higher temperatures. It was considered by the present authors that the accuracy of the data did not warrant the application of such a detailed method for their studies.

Apart from its effect on evaporation, temperature itself has a profound influence on plant growth and on the growth and reproduction of animals. No experimental work on the temperature requirements of the plants constituting the vegetative cover of the sheep-raising areas of Queensland is known. Klages (1942), discussing temperature efficiencies in relation to crop distribution, quotes a number of authorities giving a general "zero of vital temperature point" as 40°F. or 4·4°C. Forster (1941) gives 42°F. as the lowest limit for flax growth.

Lee and his co-workers (1947) found adult sheep to be remarkably tolerant of high temperatures, but plane of nutrition had a marked effect on their reactions. Gunn and his co-workers (1942) studied the effects of high temperatures on the morphology, motility and longevity of spermatozoa produced by rams enjoying normal sexual health. They found that animals subjected to daily maximum temperatures of over 95°F., under conditions of low humidity, for a period of 28 days became comparatively infertile. Recovery, as judged by the seminal picture, took place in about 2½ months if the rams were adequately fed and if the weather was cool.

Table

Values for various factors

Station.	Late Spring. October–November.					Early Summer. December–January.					Late Summer. February–March.				
	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.
Burketown	1·81	82·9	·63	20·16	·09	12·51	85·5	·48	15·36	·81	11·38	83·7	·40	12·80	·89
Cloncurry	1·68	83·8	·79	25·28	·07	7·03	87·6	·76	24·48	·29	6·65	84·9	·63	20·32	·33
Croydon	2·26	86·9	·74	23·68	·10	13·14	86·6	·51	16·32	·81	10·80	84·5	·42	13·60	·79
Donor's Hill ..	2·24	85·0	·76	24·32	·09	11·55	86·8	·61	19·52	·59	9·32	84·1	·47	15·20	·61
Normanton	2·22	85·7	·62	19·84	·11	16·77	86·3	·42	13·60	1·23	15·93	84·6	·33	10·56	1·51
Georgetown	2·48	83·1	·65	20·96	·12	13·90	83·8	·48	15·52	·90	12·45	81·6	·40	12·96	·96
Hughenden	2·07	80·8	·67	21·44	·10	7·12	84·6	·60	19·20	·37	6·04	81·5	·45	14·56	·41
Mount Surprise ..	2·57	79·2	·50	16·00	·16	12·60	81·2	·41	13·28	·95	11·61	78·2	·31	9·92	1·17
Richmond	1·80	82·4	·73	23·52	·08	6·81	86·1	·68	21·76	·31	6·42	83·1	·53	16·96	·38
Atherton	3·39	70·8	·28	9·12	·37	19·13	74·4	·23	7·36	2·60	19·41	72·4	·14	4·64	4·18
Cairns	5·98	77·9	·32	10·24	·58	25·23	81·7	·31	9·92	2·54	33·81	80·7	·24	7·84	4·31
Cooktown	3·53	80·1	·32	10·24	·34	21·00	82·1	·27	8·64	2·43	29·02	81·1	·22	7·04	4·12
Herberton	3·57	70·4	·27	8·80	·41	15·31	73·5	·23	7·52	2·04	15·72	72·1	·15	4·80	3·28
Port Douglas ..	5·92	77·2	·29	9·28	·64	24·28	80·4	·28	8·96	2·71	32·35	79·6	·24	7·68	4·21
Cardwell	6·19	75·2	·26	8·32	·74	25·07	79·5	·25	8·16	3·07	32·87	78·9	·19	6·08	5·41
Innisfail	9·59	75·2	·21	6·72	1·43	31·74	79·4	·22	7·04	4·51	49·38	78·6	·14	4·64	10·64
Townsville	3·20	78·2	·36	11·68	·28	16·32	81·3	·33	10·56	1·55	18·38	80·7	·30	9·76	1·88
Ayr	2·70	76·6	·36	11·68	·23	15·16	80·8	·34	11·04	1·37	15·60	79·7	·28	9·12	1·71
Bowen	2·27	77·4	·34	11·04	·21	14·28	81·2	·32	10·40	1·37	14·52	80·5	·27	8·80	1·65
Mackay	4·82	75·0	·29	9·28	·52	20·99	79·5	·29	9·28	2·26	23·85	78·3	·21	6·88	3·47
St. Lawrence ..	4·20	74·8	·33	10·72	·39	13·94	79·6	·33	10·56	1·32	12·96	78·7	·27	8·64	1·50
Charters Towers ..	2·17	78·5	·42	13·60	·16	8·67	81·9	·41	13·28	·65	8·20	79·5	·33	10·72	·76
Blackall	2·87	78·5	·59	19·04	·15	5·36	84·9	·67	21·60	·25	5·91	82·3	·53	16·96	·35
Clermont	3·39	77·3	·46	14·72	·23	8·63	81·9	·45	14·56	·59	7·35	78·7	·34	11·04	·67
Emerald	3·64	77·1	·44	14·08	·26	7·42	82·1	·44	14·24	·52	6·25	79·8	·36	11·52	·54
Springsure	4·07	75·7	·43	13·92	·29	7·41	8·07	·44	14·08	·53	6·64	78·5	·38	12·16	·55
Tambo	3·12	75·2	·50	16·16	·19	5·49	81·4	·57	18·24	·30	5·46	79·0	·44	14·24	·38
Barcaldine	2·53	78·2	·58	18·56	·14	5·38	84·0	·61	19·68	·27	5·52	81·4	·50	16·00	·34
Isisford	2·35	79·2	·60	19·36	·12	4·33	85·4	·65	20·80	·21	5·50	82·8	·53	17·12	·32
Longreach	2·09	79·7	·61	19·68	·11	4·09	85·5	·67	21·44	·19	5·87	82·8	·52	16·64	·35
Camooweal	1·74	83·1	·77	24·80	·07	5·73	86·9	·74	23·68	·24	5·48	84·1	·58	18·72	·29
Kynuna	1·78	81·7	·67	21·60	·08	5·49	86·3	·68	21·76	·25	5·56	83·1	·53	17·12	·32
Urandangie	1·23	81·3	·84	26·88	·05	3·53	87·5	·91	29·28	·12	3·97	84·4	·73	23·52	·17
West Leichhardt ..	2·33	81·4	·73	23·52	·10	6·75	85·6	·70	22·40	·30	5·69	82·0	·57	18·40	·31
Winton	1·92	81·5	·71	22·88	·08	4·96	86·9	·75	24·00	·21	5·38	84·0	·58	18·72	·29
Boulia	1·40	81·0	·78	25·12	·06	2·94	87·7	·88	28·16	·10	3·50	85·4	·74	23·68	·15
Windorah	1·53	78·7	·70	22·56	·07	2·72	86·3	·86	27·52	·10	3·26	83·9	·74	23·68	·14
Bundaberg	4·81	72·3	·29	9·44	·51	13·63	77·4	·32	10·24	1·33	11·71	76·5	·26	8·32	1·41
Childers	5·50	72·7	·28	9·12	·60	12·98	77·1	·28	9·12	1·42	11·34	76·0	·23	7·36	1·54
Gayndah	5·37	72·9	·35	11·20	·48	8·76	78·1	·36	11·52	·76	7·20	76·3	·31	9·92	·73
Gladstone	4·72	73·8	·31	9·92	·48	12·37	78·3	·30	9·76	1·27	12·06	77·6	·27	8·64	1·40
Mount Morgan ..	4·32	74·4	·31	9·92	·44	9·97	78·3	·30	9·76	1·02	8·63	76·6	·26	8·48	1·02
Rockhampton ..	4·22	76·4	·37	11·84	·36	12·39	80·6	·36	11·52	1·08	12·11	79·2	·29	9·44	1·28
Westwood	4·01	73·8	·32	10·24	·39	8·80	78·5	·32	10·40	·85	7·50	76·6	·28	9·12	·82
Gympie	6·02	72·0	·28	8·96	·67	11·97	77·1	·27	8·80	1·36	12·98	75·5	·21	6·88	1·89
Ipswich	5·33	72·7	·35	11·36	·47	8·29	78·1	·37	12·00	·69	8·48	76·1	·31	10·08	·84
Maryborough ..	5·97	72·7	·31	9·92	·60	12·10	77·5	·32	10·40	1·17	12·67	76·3	·24	7·84	1·62

P = mean rainfall in inches.

$$T = \frac{\text{mean maximum} + \text{mean minimum temperature °F.}}{2}$$

S.D. = saturation deficit in inches.

for 65 Queensland stations.

Autumn. April–May.					Winter. June–July.					Early Spring. August–September.					P/E. Diff.
P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	
1·22	77·1	·51	16·48	·07	·32	69·0	·41	13·12	·02	·11	73·4	·52	16·80	·01	+ ·87
1·17	75·0	·54	17·44	·07	·96	64·9	·35	11·36	·08	·42	71·4	·54	17·28	·02	+ ·25
1·34	79·2	·51	16·48	·08	·49	71·4	·40	12·80	·04	·32	77·3	·56	18·08	·02	+ ·75
1·40	76·1	·53	16·96	·08	·70	67·7	·40	12·80	·05	·30	73·9	·56	18·08	·02	+ ·56
1·93	79·6	·47	15·04	·13	·56	72·1	·35	11·36	·05	·16	77·1	·49	15·68	·01	+ 1·46
1·56	75·3	·43	13·76	·11	·74	68·2	·34	11·04	·07	·54	73·2	·47	15·04	·04	+ ·90
1·65	71·7	·37	12·00	·14	1·38	62·6	·24	7·68	·18	·71	69·1	·40	12·96	·05	+ ·23
1·96	71·3	·30	9·76	·20	·99	64·3	·22	7·20	·14	·38	69·1	·34	11·04	·03	+ 1·03
1·41	73·2	·44	14·68	·10	1·13	63·7	·29	9·44	·12	·37	70·3	·48	15·36	·02	+ ·26
6·50	66·1	·12	3·84	1·69	2·74	61·0	·10	3·20	·86	1·62	63·3	·17	5·44	·30	+ 3·32
15·80	75·5	·19	6·24	2·53	4·43	70·4	·18	5·92	·75	3·40	72·0	·24	7·68	·44	+ 3·56
11·58	77·5	·22	7·04	1·64	2·95	73·1	·19	6·08	·49	1·76	74·9	·24	7·84	·22	+ 3·64
5·51	66·1	·12	4·00	1·38	2·02	60·4	·09	3·04	·66	1·19	63·0	·18	5·76	·21	+ 2·61
11·96	75·1	·22	7·04	1·70	3·03	69·8	·19	6·24	·49	2·23	71·3	·24	7·68	·29	+ 3·72
12·37	73·0	·16	5·12	2·42	3·39	66·1	·13	4·32	·78	2·78	68·1	·16	5·28	·53	+ 3·32
32·37	73·2	·11	3·68	8·79	11·98	67·2	·10	3·20	3·74	8·43	68·9	·13	4·16	2·02	+ 6·90
4·60	75·3	·32	10·40	·44	1·98	68·7	·26	8·48	·23	1·24	71·5	·30	9·76	·13	+ 1·65
3·56	73·7	·27	8·80	·40	2·15	66·2	·21	6·72	·32	1·85	68·9	·26	8·32	·22	+ 1·39
3·95	74·0	·24	7·68	·51	2·57	66·7	·20	6·56	·39	1·43	69·7	·25	8·00	·18	+ 1·26
9·88	70·9	·17	5·60	1·76	4·40	63·1	·13	4·32	1·02	2·68	66·1	·19	6·08	·44	+ 2·45
4·51	71·2	·23	7·52	·60	3·88	63·4	·16	5·28	·73	2·04	66·5	·22	7·20	·28	+ ·77
2·26	72·2	·30	9·60	·24	1·96	64·5	·21	6·88	·28	1·31	69·2	·29	9·28	·14	+ ·48
2·71	68·5	·33	10·72	·25	2·32	57·9	·18	5·92	·39	1·40	64·4	·23	10·56	·13	− ·14
2·90	68·1	·26	8·48	·34	2·74	58·7	·16	5·12	·54	1·64	64·5	·27	8·64	·19	+ ·13
2·43	68·9	·27	8·64	·28	2·80	59·4	·17	5·44	·51	1·91	64·7	·27	8·64	·22	+ ·03
2·76	67·5	·30	9·60	·29	2·94	57·7	·19	6·24	·47	2·19	63·5	·29	9·28	·24	+ ·08
2·72	65·7	·29	9·28	·29	2·45	55·7	·15	4·96	·49	1·60	61·8	·29	9·44	·17	− ·11
2·53	69·4	·34	10·88	·23	2·15	59·4	·20	6·56	·33	1·23	65·2	·35	11·20	·11	+ ·01
2·29	69·2	·35	11·20	·20	1·96	58·9	·21	6·88	·28	1·15	65·3	·35	11·20	·10	+ ·04
1·78	70·0	·36	11·68	·15	1·68	59·6	·21	6·72	·25	·82	65·9	·36	11·68	·07	+ ·10
·79	73·4	·51	16·48	·05	·92	63·3	·33	10·72	·09	·33	70·5	·51	16·48	·02	+ ·20
1·22	72·4	·42	13·44	·09	1·25	61·9	·24	7·68	·16	·44	68·7	·41	13·12	·03	+ ·16
·86	71·2	·53	16·96	·05	·81	60·7	·32	10·40	·08	·35	67·7	·51	16·32	·02	+ ·09
·98	72·4	·47	15·04	·07	·77	62·6	·32	10·40	·08	·41	68·3	·46	14·72	·03	+ ·24
1·37	72·2	·43	13·76	·10	1·46	61·6	·26	8·32	·18	·61	68·2	·43	13·92	·04	+ ·11
·97	70·9	·47	15·04	·06	·85	60·0	·26	8·32	·10	·52	67·1	·44	14·24	·04	+ ·05
1·55	68·3	·40	12·80	·12	1·31	57·1	·21	6·72	·19	·84	64·2	·39	12·64	·07	− ·05
5·82	68·9	·20	6·56	·89	4·69	61·1	·15	4·80	·98	2·84	64·1	·20	6·40	·44	+ ·43
4·89	68·6	·19	6·24	·78	4·15	61·1	·16	5·12	·81	2·98	64·3	·21	6·72	·62	+ ·73
2·98	66·3	·23	7·36	·40	3·28	57·5	·15	4·96	·66	2·70	61·7	·23	7·36	·37	+ ·07
4·30	70·6	·25	8·00	·54	4·49	63·0	·17	5·44	·83	2·21	65·9	·23	7·36	·30	+ ·57
3·39	68·1	·21	6·88	·49	3·80	59·7	·14	4·48	·85	2·01	64·6	·21	6·88	·29	+ ·17
4·12	71·6	·24	7·84	·53	4·34	62·9	·17	5·44	·80	2·10	67·3	·24	7·68	·27	+ ·48
3·46	66·9	·22	7·20	·48	3·60	58·8	·14	4·64	·78	2·13	63·3	·21	6·88	·31	+ ·04
6·21	66·7	·15	4·80	1·29	4·75	58·1	·11	3·68	1·29	3·80	61·8	·16	5·12	·74	+ ·60
4·31	63·5	·19	6·08	·71	3·68	57·5	·14	4·64	·79	3·13	61·7	·22	7·04	·44	+ ·05
6·82	68·2	·19	6·08	1·12	4·92	60·4	·13	4·32	1·14	3·57	63·4	·20	6·56	·54	+ ·48

E = evaporation (calculated from saturation deficit) in inches.

P/E = ratio of precipitation to evaporation.

P/E Diff. = P/E for January-February-March—P/E for June-July.

Table 1

Station.	Late Spring. October–November.					Early Summer. December–January.					Late Summer. February–March.				
	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.
Dalby	4·99	71·0	·37	11·84	·42	6·86	76·6	·39	12·64	·54	5·53	74·5	·34	10·88	·51
Goondiwindi.. ..	3·95	72·3	·39	12·48	·31	5·90	79·5	·48	15·36	·38	4·90	77·2	·39	12·48	·39
Pittsworth	4·97	68·3	·31	10·08	·49	7·57	73·5	·32	10·24	·74	5·48	71·9	·29	9·28	·59
Stanthorpe	5·22	63·0	·24	7·84	·67	7·22	69·4	·28	8·96	·81	5·68	67·7	·22	7·20	·79
Wallangarra ..	5·60	62·1	·22	7·04	·80	7·81	68·0	·28	9·12	·86	5·57	66·8	·24	7·84	·71
Warwick	4·95	67·6	·28	9·12	·54	7·14	73·7	·33	10·56	·68	5·64	71·8	·27	8·80	·64
Miles	4·57	72·6	·38	12·32	·37	6·99	78·7	·42	13·60	·51	5·42	76·6	·34	10·88	·50
Mitchell	3·66	72·0	·46	14·72	·25	5·79	79·2	·53	16·96	·34	5·56	76·5	·42	13·44	·41
Roma	3·85	73·7	·45	14·40	·27	5·69	80·2	·51	16·32	·35	5·47	78·0	·42	13·60	·40
St. George	2·94	73·8	·44	14·08	·21	4·68	81·8	·56	17·92	·26	4·39	78·9	·45	14·40	·30
Surat	3·53	73·2	·38	12·32	·29	5·55	80·5	·47	15·20	·37	5·30	78·1	·39	12·48	·42
Yuleba	4·35	72·8	·40	12·80	·34	5·91	78·6	·43	13·76	·43	5·74	76·7	·37	11·84	·48
Bollon	2·65	74·2	·44	14·24	·19	4·50	82·1	·55	17·60	·26	3·59	79·7	·45	14·40	·25
Charleville	2·87	75·5	·54	17·28	·17	4·82	82·7	·65	20·64	·23	4·80	80·3	·53	16·96	·28
Cunnamulla	1·87	74·7	·55	17·76	·11	3·03	82·7	·67	21·60	·14	3·30	80·1	·55	17·60	·19
Adavale	2·13	77·0	·63	20·16	·11	3·95	84·6	·76	24·48	·16	3·82	82·5	·65	20·96	·18
Thargomindah .	1·65	76·1	·64	20·64	·08	2·74	84·5	·83	26·56	·10	2·14	82·3	·71	22·72	·09

P = mean rainfall in inches.

$$T = \frac{\text{mean maximum} + \text{mean minimum temperature °F.}}{2}$$

S.D. = saturation deficit in inches.

—continued.

Autumn. April–May.					Winter. June–July.					Early Spring. August-September.					P/E. Diff.
P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	P.	T.	S.D.	E.	P/E.	
2·65	63·1	·21	6·88	·39	3·33	53·6	·12	3·84	·87	2·78	58·7	·22	7·04	·39	— ·36
3·04	63·7	·21	6·88	·44	3·43	53·5	·11	3·52	·97	2·68	59·0	·20	6·56	·41	— ·58
2·98	61·2	·20	6·56	·45	3·52	52·1	·12	3·84	·92	2·76	57·0	·19	6·08	·45	— ·33
3·42	56·0	·13	4·16	·82	3·92	46·7	·08	2·56	1·53	3·97	51·0	·14	4·48	·88	— ·74
2·96	56·2	·15	4·96	·60	4·22	47·4	·09	3·04	1·39	3·40	51·0	·16	5·12	·66	— ·68
3·06	60·7	·16	5·28	·58	3·48	51·3	·09	2·88	1·21	3·21	55·5	·16	5·12	·63	— ·57
2·89	63·6	·21	6·88	·42	3·26	53·4	·12	4·00	·82	2·42	59·2	·22	7·20	·34	— ·31
2·56	62·3	·22	7·20	·36	2·98	52·4	·11	3·68	·81	2·13	58·1	·24	7·68	·28	— ·40
2·68	64·1	·24	7·84	·34	2·94	53·9	·12	3·84	·77	2·26	60·1	·23	7·52	·30	— ·36
2·66	64·6	·28	8·32	·29	2·70	53·8	·12	4·00	·68	1·97	59·8	·23	7·36	·27	— ·37
2·50	64·5	·23	7·52	·33	3·38	53·9	·11	3·68	·97	2·26	59·9	·21	6·88	·33	— ·50
2·73	62·9	·22	7·20	·38	3·22	52·9	·12	4·00	·81	2·20	58·8	·22	7·20	·31	— ·33
2·24	64·4	·23	7·52	·30	2·42	53·4	·11	3·68	·66	1·91	59·5	·22	7·04	·27	— ·40
2·52	65·6	·30	9·60	·26	2·51	54·7	·14	4·64	·54	1·60	61·0	·29	9·44	·17	— ·26
2·12	65·0	·29	9·44	·22	2·01	54·3	·14	4·64	·43	1·48	60·8	·29	9·28	·16	— ·25
2·01	66·8	·34	10·88	·18	2·17	55·6	·18	5·76	·38	1·13	62·3	·33	10·72	·11	— ·20
1·58	66·0	·35	11·20	·14	1·25	54·7	·17	5·60	·22	·99	61·5	·34	10·88	·09	— ·13

E = evaporation (calculated from saturation deficit) in inches.

P/E = ratio of precipitation to evaporation.

P/E Diff. = P/E for January–February–March—P/E for June–July.

DATA AVAILABLE.

Limited data were available and these consisted of:

(i.) Monthly rainfall totals for periods of 50 years and over for 65 Queensland stations. The mean was calculated and is set down with other data used in Table 1.

(ii.) The mean monthly maximum and minimum dry bulb readings and humidities for 65 stations for a period of from 20 to 30 years. Saturation deficits calculated from these figures are set out in Table 1.

Monthly maximum and minimum temperatures and humidities for individual years for Charleville, Cloncurry and Longreach.

(iii.) Records of evaporation from a free water surface from five stations, viz., Winton, Blackall, St. George, Charleville and Warwick. These are given in Table 2.

Table 2.

Relationship between saturation deficit and measured evaporation for five Queensland stations.

Station.	Value.	Jan.	Feb.	Mar.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
Winton	S.D.	·72	·61	·56	·50	·36	·26	·26	·36	·51	·67	·76	·78
	E.	12·14	9·22	9·29	7·38	5·74	4·06	4·28	5·96	8·02	10·17	10·96	12·14
	k	16·9	15·1	16·6	16·7	16·0	15·6	16·5	16·5	15·7	15·2	14·4	15·6
Blackall	S.D.	·67	·57	·49	·41	·26	·18	·19	·27	·39	·56	·63	·68
	E.	10·85	8·55	8·12	6·17	4·35	2·96	3·12	4·51	6·36	8·85	9·81	10·88
	k	16·2	15·0	16·6	15·0	16·7	16·5	16·4	16·7	16·3	15·8	15·6	16·0
Charleville	S.D.	·65	·58	·48	·37	·23	·14	·15	·24	·35	·49	·59	·64
	E.	10·70	8·80	8·18	6·13	4·19	2·65	2·71	4·00	5·89	7·99	9·21	10·29
	k	16·5	15·2	17·1	16·6	18·2	18·8	18·1	16·7	16·8	16·3	15·6	16·1
St. George	S.D.	·61	·50	·40	·31	·21	·13	·12	·19	·27	·40	·48	·51
	E.	10·29	8·79	8·02	5·35	3·52	2·33	2·34	3·46	5·46	7·90	8·93	9·94
	k	16·9	17·5	20·0	17·2	16·7	17·9	19·5	18·2	20·2	19·7	18·6	19·5
Warwick	S.D.	·32	·29	·26	·19	·14	·09	·09	·13	·19	·27	·30	·34
	E.	7·22	5·97	5·75	4·05	2·89	1·98	1·99	2·69	3·98	5·60	6·41	7·21
	k	22·5	20·6	22·1	21·3	20·6	22·0	22·0	20·7	20·9	20·7	21·3	21·2

S.D.—Saturation Deficit. E.—Measured Evaporation in inches. k—constant (in equation $E = k \times S.D.$).

METHODS.

Temperature.

Mean minimum temperatures for June-July were plotted for all stations and isolines drawn (Figure 1). Mean monthly maximum temperatures for January-February were treated similarly.

The number of months having an average maximum temperature of more than 95°F. was calculated for each station for which data were recorded and the results were plotted on the map shown in Figure 2.

Rainfall Distributions.

Histograms showing the frequency distribution of monthly rainfalls were compiled with class intervals of 0·25 inch. Coefficients of variations for mean monthly rainfalls of a number of stations were calculated.

P/E Ratios.

The lack of data, particularly of daily or weekly rainfalls, made it necessary to work on a monthly basis. With evaporation records so few and so widely distributed it was impossible to attempt any detailed mapping on the basis of recorded evaporation and some method of calculating or estimating evaporation from available data had to be evolved. Davidson's (1935) method of conversion expressed as the formula $E = k \times S.D.$ was used and, to find a value for K (mean monthly), evaporation figures for five stations were compared with values for saturation deficits calculated from records of temperature and relative humidity. In these calculations temperatures used were $\frac{\text{Max.} + \text{Min.}}{2}$ and relative humidities those at 9 a.m.

Table 2 gives a comparison of monthly evaporations and calculated saturation deficits for the five inland stations in Queensland. Values of k range from 14.9 to 21, but most of them are about 16 (coastal about 22). It was considered that in a preliminary study such as this the use of a single value of k is justified. Accordingly figures for saturation deficit were converted to evaporation figures by multiplying by 16. The chief reason for converting saturation deficit to evaporation instead of using Meyer ratio $\frac{P}{S.D.}$ was to have both precipitation and evaporation in the same unit, namely, inches.

Monthly evaporation figures for the 65 stations were computed from the calculated values for saturation deficit. Examination of the frequency distribution charts of mean monthly rainfall as well as evaporation figures showed that the months could be grouped in pairs beginning with October-November, thus dividing the year into six periods, each of two months. Bi-monthly P/E ratios were then calculated and maps for each two-monthly "season" are given as Figures 3-8.*

---

* *Marginal Errors.*—In drawing charts of a limited area, especially one with arbitrary boundaries such as Queensland, there is always the possibility of marginal error. Such error can be minimized by plotting relevant data on stations outside the region being studied. Without such data the correct direction of isolines at and near the margins cannot be determined with certainty.

In order to reduce marginal error, data were plotted for coastal stations using the same values of k as for the inland stations. It must be emphasized that the coastal data are not exhaustive and the isolines shown on the maps do not represent the complete picture. Further detailed study is needed in the agricultural regions of the coastal belt. Although they are outside the scope of the present paper, the small amount of work which has been done indicates that the methods outlined here should yield useful results if applied to the dense network of stations of the coastal regions.

Along the western and southern borders the possibility of marginal error is greater. Detailed reports from stations in New South Wales, South Australia and Northern Territory were not available. The direction of isolines along these margins is only approximate.

P/E Differences.

A review of the frequency distribution charts of mean monthly rainfall showed that in semi-arid Queensland rainfall occurs in two well-marked seasons:

(1) Summer season (January, February and March);

(2) Winter season (June and July).

Rainfall in the summer season is far greater than in winter. Falls do occur in other months—notably October, November and December—but these are usually storms of erratic distribution and rarely represent general rain. It is generally believed that winter rainfall is more reliable in the south and becomes negligible towards the north.

In an attempt to fix the northern limits of effective winter rainfall the differences between January-February-March P/E and June-July P/E ratios were plotted and isolines drawn at intervals of 0.1 (Figure 9).

South of the zero line, P/E for winter exceeds P/E for summer; north of the line summer P/E is the greater. Maps were also drawn showing the differences between the following groups of months:

(1) February-March—June-July.

(2) December-January—June-July.

In these, the lines followed the same general pattern as in Figure 4 and the position of the zero line was approximately the same.

In order to test the consistency of the general pattern for different seasons over individual years, 11-year and 23-year averages of the summer minus winter P/E values were calculated for the three centres for which these data were available.

Seasonal Distribution and Reliability of Effective Rainfall.

Examination of Queensland data in the light of known conditions in the field indicated that for summer months Trumble's (1939) figure of 0.3 (for a P/E value) as being indicative of "a period of influential rain" is too high, because good growth of pastures may take place when monthly P/E values are less than 0·3. Trumble (1939) defined "a period of influential rainfall" as one in which average P/E is equal to or greater than 0.3, while for tropical Australia Lawrence (1941) used the lower value of 0.25. It was considered that for summer months both of these figures are too high; good growth of pastures can take place during months in which average P/E is less than 0·3 or 0·25. In addition, as Trumble (1939) pointed out, Queensland rainfall is more variable than that of southern States, where the period of minimum evaporation coincides with the period of maximum precipitation. In Queensland, precipitation and evaporation figures tend to run parallel and evaporation is high during the month of maximum precipitation. Under such conditions, a lower value for P/E ratio might be adopted as the criterion of a "period of influential rain." Tentative tests of lower values indicated that for summer a wet month—i.e., month of effective rainfall—could be defined as one with a P/E equal to or greater than 0·2. Accordingly this value was used for the period October to March. In winter the figure of 0·3 was found to agree

fairly well with observed field conditions, and this was used for the months from April to September. It must be emphasized that this lower figure is purely arbitrary and has not been reached as the result of any experimental work on rate of loss of moisture from soils, but field observations indicate that it approximates the P/E value which will initiate and sustain growth of natural vegetation.

Except for Cloncurry, Longreach and Charleville, data for saturation deficit were available only in the form of monthly means for the whole period on record. This made it impossible to calculate individually the actual P/E ratio for each month. It was believed that, as variability of evaporation was small compared with that of rainfall, it might be possible to use average values for evaporation combined with individual figures for rainfall without introducing any very great error.

To test this, the number of wet months was determined for the three stations, using (1) ratio of precipitation to evaporation, the latter being obtained by calculation from actual values for saturation deficit for each month; and (2) ratio of precipitation to evaporation, in which evaporation figures were calculated from mean values of saturation deficit for the whole period on record.

Table 3.

Comparison of two methods of computing P/E.

Percentage of Years in which Months Receive an Effective Rainfall.

Station.	Oct.–Nov.		Dec.–Jan.		Feb.–Mar.		Apr.–May.		June–July.		Aug.–Sept.	
	Ind. E.*	Av. E.†	Ind. E.	Av. E.	Ind. E.	Av. E.	Ind. E.	Av. E.	Ind. E.	Av. E.	Ind. E.	Av. E.
Cloncurry	16	8	50	54	66	66	12	4	20	12	0	0
Longreach	21	19	47	41	52	55	27	18	42	42	6	6
Charleville	38	41	60	63	51	51	25	25	70	70	28	28

* Ind. E. — Percentage of years in which months receive an effective rainfall, taking evaporations for individual years.

† Av. E. — Percentage of years in which months receive an effective rainfall, taking a mean evaporation figure for all years.

The comparison is set out in Table 3. It shows that the mean saturation deficit may be used for computing evaporation without materially affecting the results.

Zonation.

Zonation was based on the three factors which were considered to be most important in the study of a pastoral environment, viz:—(1) effective rainfall; (2) distribution of rainfall; (3) reliability of rainfall. In addition it was felt that where necessary consideration could be given to the mean annual rainfall.

From the distribution and reliability studies Figures 10, 11 and 12, showing the number of wet months in summer and winter in 50, 66 and 75 per cent. respectively of all years recorded, were drawn.

Figure 1.

Map showing mean minimum temperatures for the months of June and July.

Areas having 6, 4, 2 or 0 wet summer months in 50 per cent. of the years are given the indices $\frac{A}{50}$, $\frac{B}{50}$, $\frac{C}{50}$ and $\frac{D}{50}$, while for winter a, b, c, and d, are used in a similar manner. An area receiving effective rain in six summer months and six winter months (though not necessarily in the same year) in 50 per cent. of years studied is denoted by $\frac{Aa}{50}$. A zone having four months of effective rain in the summer and two months of effective rain during the winter in 50 per cent. of the years is denoted by $\frac{Bc}{50}$.

Likewise, on the map showing the conditions for 66 per cent. and 75 per cent. of the years, the index used for an area having six summer and four winter months during which effective rain fell is classified as $\frac{Ab}{66}$ or $\frac{Ab}{75}$.

At the same time consideration has been given, where necessary, to the mean annual rainfall, by prefixing this figure to the index classification. For example, Camooweal and Isisford are both in the zone $\frac{Cd}{66}$, but the mean annual rainfall at the former is 14.89 inches and at the latter 17.85 inches. These centres can be differentiated as $15\frac{Cd}{66}$ for Camooweal and $18\frac{Cd}{66}$ for Isisford.

Table 4 shows, for each of the stations considered, the zonation index for each order of reliability, as well as mean annual rainfall, mean maximum summer and minimum winter temperatures, and number of months having a mean maximum temperature equal to or greater than 95°F.

RESULTS.

Temperature.

The map showing mean minimum temperatures for June and July (Figure 1) is characterized by isolines which, in the inland area, run more or less east-west. In the east they are modified by ranges and proximity of the sea. Around Mitchell, Yuleba and Surat is a peculiar cold closed zone within the 40°F. isotherm.

It is considered that Queensland maximum summer temperatures are not at any time high enough to inhibit appreciably the growth of plants and consequently the map showing these isolines is not published.

Figure 2, showing the number of months having a mean maximum temperature of more than 95°F., exhibits a closed area around Kynuna, Richmond and Winton in which six months exceed the above value. The zone which has five hot months is of somewhat greater dimensions, extending from Croydon in the north to a point west of Longreach in the south, and including Donor's Hill, Cloncurry, Camooweal, Urandangie, and Boulia.

The line forming the boundary of the zone, which has four hot months, runs more or less east from the border north of Camooweal, turning south near Croydon, and again assuming an east-west direction from Blackall to the South Australian border. It includes the districts of Hughenden, Longreach, Isisford, Blackall and Windorah.

Figure 2.

Map showing the mean number of months with mean maximum temperature 95°F. or greater.

The western limits of the 6-months line are somewhat indefinite. Camooweal, Cloncurry and Urandangie have a sixth month in which the mean maximum temperatures are 94·8°, 94·6° and 94·7° respectively.

The lines bounding the other zones follow approximately the same pattern with a progressively decreasing tendency to run east-west in the southern districts.

The area in which the mean maximum temperature does not exceed 95°F. in any month extends as far west as Mt. Surprise and Charters Towers in the north, while towards the south it includes the Springsure, Mitchell, Surat and Goondiwindi districts.

Rainfall Distribution.

The coefficients of variations for the mean monthly rainfalls were of the order of 0·9 and accordingly were considered unsatisfactory. The histograms are not reproduced here but they were studied in detail before commencing work on the seasonal distribution and reliability of effective rainfall.

Table 4.

Zonation indices and other data for 65 Queensland stations.

Station.	Ind. 50% *	Ind. 66% †	Ind. 75% ‡	Mean Annual Rainfall. Inches.	Jan.–Feb. Mean Max. Temp. °F.	June–July Mean Min. Temp. °F.	No. of Months with Mean Max. T. > 95° F.
Burketown	Bd	Bd	Bd	27·61	92·9	56·3	1
Cloncurry	Cd	Cd	Cd	17·96	97·8	52·8	5
Croydon	Bd	Bd	Bd	28·75	95·7	58·0	5
Donor's Hill	Bd	Bd	Bd	25·49	96·2	53·7	5
Normanton	Bd	Bd	Bd	37·87	93·8	59·7	3
Georgetown	Bd	Bd	Bd	31·74	92·8	54·3	3
Hughenden	Bd	Bd	Cd	19·05	96·2	48·0	4
Mount Surprise ..	Bd	Bd	Bd	30·28	90·7	49·7	0
Richmond	Bd	Cd	Cd	18·08	97·6	48·6	6
Atherton	Aa	Bb	Bb	53·75	83·2	50·8	0
Herberton	Ab	Ab	Ab	43·80	81·6	50·2	0
Cairns	Aa	Ab	Ab	88·75	89·5	62·4	0
Cooktown	Ab	Bb	Bb	69·32	88·6	67·0	0
Port Douglas ..	Ab	Ab	Ab	79·66	86·2	63·4	0
Cardwell	Aa	Ab	Ab	82·38	87·7	56·0	0
Innisfail	Aa	Aa	Aa	143·78	87·4	58·5	0
Townsville	Bc	Bc	Bd	45·72	87·1	60·8	0
Ayr	Bc	Bc	Bc	41·54	89·2	54·7	0
Bowen	Bc	Bc	Bc	39·30	87·8	58·0	0
Mackay	Ab	Ab	Bb	66·91	85·7	54·7	0
St. Lawrence ..	Ab	Ab	Ac	41·11	87·9	52·5	0
Charters Towers ..	Bc	Bd	Bd	24·64	91·7	52·8	0
Blackall	Bc	Cc	Cd	20·77	98·1	44·6	4
Clermont	Ac	Bc	Bc	26·97	92·5	44·4	1
Emerald	Bc	Bc	Bc	24·61	93·6	45·7	1
Springsure	Ac	Bc	Bc	26·19	92·3	44·3	0
Tambo	Bc	Bc	Cc	21·00	95·0	40·5	1
Barcaldine	Bc	Cc	Cd	19·70	95·4	46·5	2
Isisford	Cc	Cd	Cd	17·73	98·2	45·4	4
Longreach	Cc	Cd	Cd	16·52	98·3	45·4	4

Table 4—continued.

Zonation indices and other data for 65 Queensland stations—*continued.*

Station.	Ind. 50% *	Ind. 66% †	Ind. 75% ‡	Mean Annual Rainfall. Inches.	Jan.–Feb. Mean Max. Temp. °F.	June–July Mean Min. Temp. °F.	No. of Months with Mean Max. T. > 95° F.
Camooweal	Cd	Cd	Cd	14·96	97·7	48·9	5
Kynuna	Cd	Cd	Cd	15·71	99·1	46·9	6
Urandangie	Dd	Dd	Dd	10·64	100·1	46·4	5
Winton	Cd	Cd	Cd	15·90	99·4	47·2	6
Boulia	Dd	Dd	Dd	10·34	100·7	46·8	5
Windorah	Dd	Dd	Dd	11·26	99·7	44·2	4
Bundaberg	Aa	Ab	Ab	43·47	85·8	50·3	0
Childers	Aa	Ab	Ab	41·78	87·1	51·0	0
Gayndah	Ab	Ab	Ac	30·44	89·8	43·0	0
Gladstone	Ab	Ab	Ac	39·98	85·3	53·7	0
Mount Morgan ..	Ab	Ab	Ac	32·19	87·6	48·7	0
Rockhampton ..	Ab	Ac	Ac	38·95	88·8	52·1	0
Westwood	Ab	Ac	Ac	29·32	89·2	45·6	0
Brisbane	Aa	Aa	Aa	44·72	84·9	50·0	0
Gympie	Aa	Aa	Ab	45·42	87·7	44·6	0
Ipswich	Ab	Ab	Ab	33·07	89·3	45·1	0
Maryborough ..	Aa	Ab	Ab	45·63	86·8	49·1	0
Dalby	Ab	Ab	Ab	26·12	89·0	40·8	0
Goondiwindi ..	Bb	Bb	Bb	24·05	92·8	42·1	0
Pittsworth	Ab	Ab	Ab	27·22	85·3	42·2	0
Stanthorpe	Aa	Aa	Aa	29·52	81·0	34·9	0
Wallangarra	Aa	Aa	Aa	29·48	79·3	37·6	0
Warwick	Aa	Ab	Ab	27·49	85·2	38·9	0
Killarney	Aa	Aa	Aa	28·55	84·1	37·0	0
Miles	Ab	Bb	Bc	25·63	91·5	39·6	0
Mitchell	Bb	Bc	Bc	23·00	91·9	38·6	0
Roma	Bb	Bc	Bc	23·08	93·3	40·2	0
St. George	Bb	Bc	Cc	19·65	95·2	41·6	1
Surat	Bb	Bb	Bc	22·69	93·9	40·7	0
Yuleba	Ab	Bb	Bc	24·47	92·1	37·0	0
Bollon	Cc	Cc	Cc	17·62	95·6	40·8	2
Charleville	Bb	Cc	Cc	19·38	96·2	41·2	3
Cunnamulla	Cc	Dc	Dc	14·11	96·1	42·9	2
Adavale	Cc	Dd	Dd	15·38	98·0	42·5	3
Thargomindah ..	Dd	Dd	Dd	10·56	97·1	43·7	3

* Ind. 50% — Zone in which station is situated on map showing the number of wet months in 50% of years on record.

† Ind. 66% — Zone in which station is situated on map showing the number of wet months in 66% of years on record.

‡ Ind. 75% — Zone in which station is situated on map showing the number of wet months in 75% of years on record.

Precipitation-Evaporation Ratios.

Figures 3-8, drawn from the rainfall evaporation ratios for the six two-monthly seasons, presented the following features:—

October-November: The late spring is normally a dry season, so the isolines are few and well spaced. They trend roughly north-south.

December-January: Early summer shows a marked change in both number and direction of the isolines. In this season the isolines are roughly parallel to the coastline.

February-March: The isolines in the main summer rainfall season are very similar to those of December-January, except that the lines are further towards the south-west.

April-May: The autumn season shows another marked change in the configuration of the lines. The lines are few and well spaced. A curious feature is the dry tongue which extends into the Lockyer and Fassifern Valleys (west of Brisbane) from the north-west. The general trend of the lines is north-south.

June-July: The isolines in the winter map are more numerous and more closely spaced. This is a reflection of the higher rainfall normally experienced in this season. The dry tongue apparent on the April-May map is well developed and is accentuated by a moist area further west. This extends from Warwick to Mitchell. A study of the synoptic weather charts for several years indicates that during winter cold fronts are frequently retarded east of Charleville. This area of retardation coincides very closely with the position of the moist tongue shown on the June-July P/E map. The general trend of the lines is north-south.

August-September: In early spring the pattern of lines is similar to that of October-November. Some evidence remains of the dry tongue noted on the two previous maps but almost all the lines trend north-south.

P/E Difference Map.

The zero line on the summer-winter P/E map extends through Isisford towards Barcaldine, then east almost to Emerald. From Emerald it dips sharply southward to about latitude 25.5° S., then swings eastward to Taroom. From Taroom it extends in a south-easterly direction to about Ipswich.

North and east of the zero line, P/E for summer is greater than P/E for winter; south and west of the line the reverse applies.

The moist tongue in southern Queensland noted in the June-July P/E map is reflected in the P/E difference map.

Figure 3.

Map showing isolines of P/E values for the two-monthly period October-November.

Figure 4.

Map showing isolines of P/E values for the two-monthly period December-January.



Figure 5.

Map showing isolines of P/E values for the two-monthly period February-March.

Figure 6.

Map showing isolines of P/E values for the two-monthly period April-May.



Figure 7.

Map showing isolines of P/E values for the two-monthly period June-July.

Figure 8.

Map showing isolines of P/E values for the two-monthly period August-September.

As might be expected, the low rainfall reliability did not permit the isolines shown in Figure 9 to apply to each individual year; in fact, in some years a marked departure was shown. The 11-year means showed closer agreement and the 23-year means gave a reasonably close correlation.

For Charleville, Longreach and Cloncurry, for instance, where the mean summer—winter P/E values are — ·29, + ·07 and + ·26 respectively, the number of 23-year periods with various mean values are as shown below:—

Charleville.		Longreach.		Cloncurry.	
Mean Value.	No. of Periods.	Mean Value.	No. of Periods.	Mean Value.	No. of Periods.
—·23	2	+·04	1	+·23	1
—·27	2	+·05	1	+·24	4
—·28	1	+·06	3	+·25	2
—·29	(mean) 4	+·07	(mean) 1	+·26	(mean) 4
—·30	1	+·08	1	+·29	1
—·31	1	+·09	1		
—·32	2	+·10	1		
—·35	1	+·11	1		
—·37	1	+·13	3		

SEASONAL DISTRIBUTION AND RELIABILITY OF EFFECTIVE RAINFALL.

From a study of seasonal distribution and reliability of periods of effective rain it was found possible to divide pastoral Queensland into a number of zones. These are shown in Figures 10, 11 and 12. Rainfall of at least 66 per cent. reliability is considered to be a minimum requirement for production from sheep under pastoral conditions.

The boundaries of all zones are approximate. Where a boundary line passes between two recording stations it is possible to place the line anywhere between them. In some cases—for example, in the eastern boundary of zone "Dd"—this may introduce a possible error of ± 100 miles. Although boundaries are shown on the maps as lines they are actually broad transition strips.

Zone Dd.

The reliability of the effective rain for any of the two-monthly periods into which the year was divided is less than 66 per cent. for the zone $\frac{Dd}{66}$. The eastern boundary of the zone extends from south of Camooweal through Boulia, Jundah and Adavale districts into northern New South Wales.

The eastern boundaries of these zones when considered on the 75 per cent. and 50 per cent. reliability levels follow approximately the same course as $\frac{Dd}{66}$.



Figure 9.

Map showing difference of P/E values for periods January-February-March and June-July.

Figure 10.

Map showing zonation based on the number of wet summer and winter months in 50% of the years on record.

A and a indicate 6 wet summer and 6 wet winter months respectively.
B and b indicate 4 wet summer and 4 wet winter months respectively.
C and c indicate 2 wet summer and 2 wet winter months respectively.
D and d indicate 0 wet summer and 0 wet winter months respectively.

Figure 11.

Map showing zonation based on the number of wet summer and winter months in 66% of the years on record. For legend see Figure 10.

Figure 12.

Map showing zonation based on the number of wet summer and winter months in 75% of the years on record. For legend see Figure 10.

Zone Dc.

Zone $\frac{Dc}{66}$, which extends from the Cunnamulla district northwards towards Adavale, is an area remarkable in that it is the only zone in Queensland where the period of effective winter rains is more reliable than the period of effective summer rains.

When considered on a basis of 75 per cent. reliability it is seen that the apex of this triangle retreats to the south, but on a 50 per cent. reliability the importance of the summer rains increases and some of the area is included in the zone $\frac{Cc}{50}$.

Zone Cd.

Zone $\frac{Cd}{66}$ includes Camooweal, Cloncurry, Richmond, Kynuna, Winton, Longreach, Isisford, and part of the Adavale districts.

When considered on a 75 per cent. basis this zone extends eastwards to include Hughenden, Barcaldine and Blackall. . On a basis of 50 per cent. reliability the zone is narrower and includes only Camooweal, Cloncurry, Kynuna and Winton districts.

Zone Cd runs almost parallel to the coastline. South-east of about Barcaldine-Isisford it merges into Zone Cc.

Zone Cc.

Zone $\frac{Cc}{66}$ is long and narrow and continues in a direction parallel to the coastline. It includes the Blackall-Barcaldine districts and extends southwards to Charleville and Bollon.

This zone is still important when considered on a basis of 75 per cent. reliability; it commences further south, between Blackall and Tambo, and extends further east to St. George. On a basis of 50 per cent. reliability it includes Longreach, Adavale and Cunnamulla.

Zone Bd.

Zone $\frac{Bd}{66}$ is broad in the north, where it extends from Burketown to Mt. Surprise. It narrows southward, but includes Hughenden and Charters Towers and extends on to the Alice Tableland and Lake Galilee. The whole of this zone lies north of the Tropic of Capricorn.

On a basis of 75 per cent. reliability this zone extends from the Gulf of Carpentaria to the east coast of Townsville and then narrows southwards to a point west of Emerald. On a basis of 50 per cent. reliability the zone includes the whole of the area around the Gulf of Carpentaria, extends in a south-easterly direction to include Richmond and Hughenden, and narrows to a point about Aramac (42 miles north of Barcaldine).

Zone Bc.

Zone $\frac{Bc}{66}$ extends southwards from the coast in the Townsville-Bowen area. It includes Clermont, Emerald and Springsure, the greater part of the Dawson Valley, and the towns of Mitchell, Roma, Surat and St. George. When considered on a basis of 75 per cent. reliability this zone is narrower in the latitude of Springsure, but it broadens to the south to include Roma, Yuleba and Miles. South of Surat it narrows again as it approaches the New South Wales border. Taken at the 50 per cent. reliability level, the zone includes country as far west as, and to the north of, Charters Towers. To the east it extends to the coast in the Ayr-Bowen districts, then southwards and westwards to Barcaldine, Blackall and Tambo; it excludes Clermont and Springsure but includes Emerald. South of Tambo it merges into zone $\frac{Bb}{66}$.

Zone Bb.

Zone $\frac{Bb}{66}$ extends southwards from a narrow peak about Mount Morgan, and broadens to include Yuleba, Surat, Miles and Goondiwindi. When judged on a basis of 75 per cent. reliability, this zone includes only a small area of the south-western Darling Downs and another isolated small area about Mackay. When considered on a basis of 50 per cent. effective rainfall reliability the zone includes St. George, Mitchell, Roma and Charleville.

Zone Ac.

Zone $\frac{Ac}{66}$ embraces only a small area from Rockhampton westward. When considered from the point of view of 75 per cent. reliability this area extends from St. Lawrence almost to Dalby and includes Rockhampton, Westwood and Gayndah. When judged on a basis of 50 per cent. reliability the Ac zone is located further west and embraces most of the Clermont and Springsure districts.

Zone Ab.

Zone $\frac{Ab}{66}$ covers most of the agricultural regions of the south-eastern portion of the State, and extends westward to Dalby, Pittsworth and Warwick. A small isolated area about Mackay and St. Lawrence is also included in this category.

This zone, when considered on a basis of 75 per cent. reliability, is smaller and more restricted, but still includes Dalby and Pittsworth, while for 50 per cent. reliability it extends further west and embraces Yuleba and Miles.



Figure 13.
Soil map of western Queensland, after Blake (1938).

DISCUSSION.

Soils and Vegetation in Relation to Climatic Factors Studied.

Prescott (1931) considered that "in Australia, there is abundant evidence that the seasonal rain factor is important in determining the soil type, characteristic soils being developed in summer and winter rainfall zones . . ." He also states that "in Queensland, the coastal plain soils developed under conditions of high rainfall, with readily permeable soils, are podsolic in character, except where red loams are developed under tropical rain forests—proceeding inland, black soils under savannah woodlands or open savannah are developed, while in the semi-arid zone browns and grey soils carrying open grassland or shrub steppe association, alternating with mulga, is to be observed reaching to the desert sandhills in the extreme west." Prescott also points out "that so far as Eastern Australia is concerned, most of the effect of a rise in temperature can be explained in terms of increased evaporation."

In the region to which these climatological studies apply, a close correlation between detailed soil type and climate is not obvious (see Figs. 13 and 14). Specific types appear to be determined largely by parent rock and subsoil drainage.

The accompanying vegetation map (Figure 14) is based on that of Blake (1938) with modifications from some of his unpublished data. The general pattern of vegetation types (grassland, woodland, &c.) depends primarily upon soil differences, though climate influences the floristic composition of the major plant formations. The distribution of individual species in relation to climate is also of interest. Mitchell grasses are dominant over large areas of brown gravelly and grey non-gravelly soils, the main species in some areas being curly Mitchell grass (*Astrebla lappacea*) and in others barley Mitchell grass (*Astrebla pectinata*). At present, the reasons for the differences in dominance of the two species of *Astrebla* are not clear. In general, barley Mitchell grass is more plentiful in the more arid regions but differences in soil also appear to play a part. The main Mitchell grass associations are to be found in climatic zone $\frac{Cd}{66}$. Other important Mitchell grass areas do occur, particularly at Dirranbandi in zone $\frac{Bb}{66}$, and on the old flood plain of the Warrego River extending southwards from Wyandra (in climatic zone $\frac{Dc}{66}$).

Climatic influence appears to be responsible for the gradual diminution in area and stature of the gidyea (*Acacia cambagei*) scrubs towards the western boundary of the Mitchell grassland. In the eastern part, such as around Blackall, gidyea occupies more land than Mitchell grass; in the western part, Mitchell grassland carries only scattered patches of gidyea. This species grows mainly within the zone $\frac{Cd}{66}$ and the northern part of zone $\frac{Cc}{66}$. Elsewhere it occurs in small communities, chiefly in depressions and along watercourses.



Figure 14.

Map of Queensland showing the western plant formations and major associations, from Blake (1938), with modifications from one of his unpublished maps.

Dense scrubs, somewhat similar to those in which gidyea occurs, are also formed by other Acacias, notably brigalow (*A. harpophylla*). Brigalow and gidyea overlap in many areas, but there seems to be a climatic check on the spread of the former towards the north-west and of the latter towards the south-east. Outside the zones $\frac{Bc}{66}$, $\frac{Bb}{66}$, and $\frac{Ab}{66}$, brigalow does not form dense scrubs, but occurs in patches, usually associated with gidyea. It thrives only in zones with at least two months' effective winter rainfall and reaches its maximum development in the zones with four months' effective winter rainfall. The distribution of this species is shown in Figure 15.

Despite the existence of suitable soils in the area now occupied by open eucalyptus forest, there is little development of mulga (*Acacia aneura*) east and north of zone $\frac{Ce}{66}$. The main areas of mulga in Queensland lie within zones $\frac{Dd}{66}$, $\frac{De}{66}$, and $\frac{Ce}{66}$. In zone $\frac{Dd}{66}$ its northward limit approximately coincides with the zero line on the P/E difference map. This indicates that, though mulga is extremely drought resistant, some winter rain is necessary for its best development. The distribution of mulga is given in Figure 16; while this species is shown in zone $\frac{Cd}{66}$, it occurs there only in small communities or as scattered individuals.

Two other Acacias—myall (*A. pendula*) and boree (*A. cana*)—are of particular interest. They grow in the same kind of soil and each occurs mainly as a fringe along the margins of scrub formations. The present known distribution of these trees in Queensland is shown in Figure 17. Myall extends as far to the north-west as Blackall and occurs in zones $\frac{Ce}{66}$, $\frac{De}{66}$, $\frac{Be}{66}$ and $\frac{Bb}{66}$. It does not extend north of the zero line on the P/E difference map (Figure 9). Apparently it must have at least two months' effective rainfall in winter. Boree, on the other hand, grows mainly in the zone $\frac{Cd}{66}$ to the north and west of Blackall. It extends into the northernmost portion of $\frac{Ce}{66}$ and a narrow tongue runs southward along the eastern edge of the zone $\frac{Dd}{66}$, but many of the trees in this zone are stunted. The meeting of these two species on the zero line of the P/E difference map appears to be significant.

Another interesting plant whose distribution appears to be influenced largely by climatic conditions is limebush (*Eremocitrus glauca*), heavy stands of which occur in zones $\frac{De}{66}$, $\frac{Ce}{66}$, $\frac{Be}{66}$ and $\frac{Bb}{66}$ (Figure 18). Three small isolated patches are known in zone $\frac{Cd}{66}$, and it is reported that these consist of stunted shrubs, which rarely produce fruit. In zone $\frac{Bb}{66}$ the plant is spreading so rapidly that it is regarded as a pest of pastoral lands.



Figure 15.

Map showing the distribution of brigalow (*Acacia harpophylla*) forests. Outside the shaded area brigalow occurs as isolated clumps, usually in depressions or mixed with other species.



Figure 16.

Map showing the distribution of mulga (*Acacia aneura*).



Figure 17.

Map showing the distribution of myall (*Acacia pendula*) and boree (*A. cana*).



Figure 18.

Map showing the distribution of limebush (*Eremocitrus glauca*).

Seasonal distribution of rainfall obviously controls plant growth during the year, and from field observations it is known that certain plants grow in particular seasons and that different years produce a characteristic pattern of herbage. Unfortunately, there are no data on the temperature required to initiate or inhibit growth in any of the plant species native to the area under consideration, but it has been stated frequently by pastoralists in the Roma district that there the plants respond poorly to winter rains. In view of the suggestions of Klages (1942) and Forster (1941) this may be explained by the closed area shown on Figure 1, in which the average minimum winter temperatures fall below 40°F. This is probably low enough to restrict the growth of most plants.

Besides its effect on the natural flora of semi-arid Queensland, climate also appears to influence the spread of the fauna. The zero line on the P/E difference map (Figure 9) is approximately as far north as rabbits have spread, and it appears that they are unlikely to invade those regions where no reliable winter rains occur, despite the occurrence of soil suitable for their burrows.

These observations focus attention on the significance of the zero line on the P/E difference maps.

Animal Husbandry in Relation to Climatic Factors Studied.

A review of the pastoral industry in Queensland shows that sheep-raising is carried out in the climatic zones $\frac{Dd}{66}$, $\frac{Dc}{66}$, $\frac{Cd}{66}$, $\frac{Cc}{66}$, $\frac{Bd}{66}$, $\frac{Bc}{66}$, and $\frac{Bb}{66}$. Except for the Boulia district, the eastern boundaries of the zone $\frac{Dd}{66}$ represent the western boundary to which sheep-grazing has extended. The presence of sheep in the Boulia area is of considerable interest, as their maintenance is largely dependent upon the channel associations of the braided courses of the Burke and Hamilton Rivers. In the south-west the Bulloo and Paroo Rivers play a similar, though not so important, role to the northern streams. Studies in climatology such as those described in this paper cannot take account of the effects in certain areas of what are virtually immense natural irrigation systems. It is also well known that topography, particularly the occurrence of mountain ranges, influences sheep distribution.

The northern and eastern boundaries of sheep-raising are not so clearly marked. Figures 11 and 19 indicate that sheep do not extend into the north-western part of zone $\frac{Cd}{66}$ nor into the northern, north-eastern and eastern part of zone $\frac{Bd}{66}$. The sheep which are depastured in zone $\frac{Bc}{66}$ are confined to the Clermont-Emerald-Springsure area and south of an east-west line running about 60 miles north of Roma. This line extends eastward into zone $\frac{Bb}{66}$. The eastern margin of the zone $\frac{Bb}{66}$ approximates the eastern boundary of sheep-raising under pastoral conditions, except for the Stanthorpe area, where the topography and soil type render large areas unsuitable for agriculture.



Figure 19.

Map showing the distribution of sheep in Queensland. Each dot represents 10,000 sheep. The figures are calculated on a mean maximum sheep population of a little over 20 million.

A detailed examination of the methods of sheep husbandry practised reveals some degree of stratification within the industry. Certain districts are used for the running of dry sheep while others are utilized essentially as breeding areas. Several factors have been of importance in influencing the way in which the various districts are being utilized. They may be enumerated as follows:—

(1) Climatic factors, which include total rainfall, its distribution and reliability, and extremes of temperature.

(2) Pasture types, which include type and extent of vegetation and the occurrence of poisonous plants or of "seed" or burrs likely to be harmful to wool.

(3) Topography of the country, which influences the ease with which dog-proof fences can be erected and water facilities arranged.

Obviously climate has an indirect influence on pasture types and soil formation. The components of climate, in conjunction with pasture and soil types, control the plane of nutrition of the sheep. This is probably the most important single factor to consider in sheep husbandry, as it influences wool production as well as lambing percentages.

A survey of the Queensland clip shows the occurrence of several different wool types, which are produced largely as the result of environmental differences and their effect on the plane of nutrition. While it is considered that these might be studied more fully, it is worth recording that there is a line of demarcation between different zones, classified on wool type, which follows fairly closely along the zero line of the P/E difference maps.

Average lambing percentages recorded for the whole of Queensland have never been spectacularly high. Dissection on a district basis, however, shows that there are wide variations between seasons and from district to district. Generally speaking, lambing percentages, judged on lambs marked to ewes joined, are poorer in the north-west, and are particularly bad in the Richmond-Winton-Kynuna area.

Several factors are probably of importance in influencing the low reproductive rate of Merino sheep in Queensland, which incidentally has a considerable influence on the efficiency with which wool is produced.

In the north-west the low reliability of the summer rains and the non-existence of any reliable winter rains make is desirable to join the rams only after the summer rains have fallen. This means mating at a time when the majority of rams have been subjected to a prolonged deficiency of both vitamin A and protein, and when the temperatures have been high enough to render the rams comparatively infertile. Accordingly, though the majority of ewes may have come on heat during the mating period, the chances of conception would have been low. At the same time those ewes which did conceive would be lambing on a falling plane of nutrition and at a time when surface water was failing. These conditions are not optimum for the survival

of young lambs or their mothers. On the other hand, mating in expectation of summer rains to give the ewes a high plane of nutrition during the terminal stages of their pregnancy means gambling on the occurrence of rains known to be unreliable and joining at a time when the majority of ewes are not likely to experience oestrus.

A detailed study of the effects of heat on baby lambs has not been undertaken. From field observations it is clear that the chances of survival are slender for lambs dropped during mid-summer, when conditions are hot and water and feed are so scarce as to necessitate the ewes travelling long distances in an area where shade trees are virtually non-existent. There is no doubt that pregnancy toxaemia causes heavy losses of pregnant ewes subjected to such conditions. There is also evidence to suggest that weak ewes are more likely to suffer from dystokia.

The closed pocket in the Winton-Richmond-Kynuna area (Figure 2), in which six months of each year have a mean maximum temperature of over 95°F., is within the zone $\frac{Cd}{66}$ and these factors make it particularly difficult to achieve continued success in the breeding of sheep. A study of this map shows that an area which has five months with a mean maximum monthly temperature of over 95°F. includes Boulia, Urandangie and Donor's Hill, and extends between Richmond and Hughenden and between Longreach and Winton. High lambing percentages are not recorded consistently from the districts enclosed within this line. The close proximity of the 5-, 4-, and 3-month isolines near Longreach indicates some improvement in conditions for lambing to the east of that centre. Consideration of the area to the south of the zero line in the P/E difference map (Figure 9) reveals the better lambing country in Queensland. With the equitable distribution of rainfall and moderate temperatures, this area is in a more favoured position.

On the other hand, in some of these districts, particularly those included in zones $\frac{Bd}{66}$ and $\frac{Be}{66}$ whose vegetation is of the type found mainly in or on open forest grazing country, and the broad transitional strip between $\frac{Cc}{66}$ and $\frac{Be}{66}$, into which vegetation of this type extends, the rate of replacement of ewes from natural increase is insufficient to meet the normal losses. In the Clermont, Emerald and Springsure districts, which are located in zone $\frac{Be}{66}$, a similar position exists. In these districts the losses as the result of worm infestation are considerably higher than those in the more arid areas of zone $\frac{De}{66}$, $\frac{Cd}{66}$, and the pastures do not ensure a uniformly high plane of nutrition for the sheep.

The effect of climate on the distribution and seasonal incidence of worms has been studied by Gordon (1947), who has extended his observations to include Queensland. It has been established clearly that amount and distribution of rainfall as well as temperature influence the species, extent

and seasonal occurrence of parasitic worms affecting sheep. It is also established that young sheep are more susceptible than older animals and losses are heavier and more frequent in the younger age group. The heavier loss in this group is sufficient to have an adverse effect on the rate at which the breeding portion of the flocks is likely to be replaced.

No attempt has been made to study drought, though it is considered that the methods employed in this work may be applicable to this problem.

ACKNOWLEDGEMENTS.

The authors are indebted to Mr. S. T. Blake for permission to use some of his unpublished work, to the Commonwealth Meteorological Bureau for furnishing fundamental data, and to all others who so willingly made suggestions and lent their assistance in carrying out this work.

REFERENCES.

ANDREWS, J., and MAZE, W. H. 1933. Some climatological aspects of aridity in their application to Australia. Proc. Linn. Soc. N.S.W. 58: 105-120.

BLAKE, S. T. 1938. The plant communities of western Queensland and their relationship, with special reference to the grazing industry. Proc. Roy. Soc. Qld. XLIX: 156-204.

DAVIDSON, J. 1933. The distribution of *Smynthurus viridis* L. (Collembola) in South Australia based on rainfall, evaporation and temperature. Aust. J. Exptl. Biol. Med. Sci. XI: 59-66.

————. 1934. The monthly precipitation-evaporation ratio in Australia, as determined by saturation deficit. Trans. Proc. Roy. Soc. S.Aust. LVIII: 33-36.

————. 1935. Climate in relation to insect ecology in Australia. 2. Mean monthly temperatures and precipitation: evaporation ratio. Trans. Proc. Roy. Soc. S.Aust. LIX: 107-112.

FORSTER, H. C. 1941. The use of climatic graphs in determining suitable areas for flax production. J. Dept. Agric. Vic. XXXIX: 515-524.

GORDON, H. McL. 1947. The epidemiology of parasitic diseases, with special referenee to studies with nematode parasites of sheep. Aust. Vet. Ass. 24th Ann. Conf. (Unpublished).

GUNN, R. M. C. 1942. Studies in fertility in sheep. (2) Seminal changes affecting fertility in rams. Counc. Sci. Industr. Res. (Aust.) Bull. 148.

KLAGES, K. H. W. 1942. Ecological Crop Geography. New York: MacMillan and Co.

LAWRENCE, E. F. 1937. A climatic analysis of New South Wales. Aust. Geogr. III: 1-23.

————. 1941. Climatic regions in tropical Australia. Aust. Geogr. IV: 20-26.

LEE, D. K. H., and ROBINSON, K. W. 1941. Reactions of the sheep to hot atmospheres. Proc. Roy. Soc. Qld. LIII: 189-200.

————. 1947. The effect of the nutritional plane upon the reactions of animals to heat. J. Ani. Sci. 6: 182-194.

MOREAU, R. E. 1938. Climatic classification from the standpoint of East African biology. J. Ecol. XXVI:

PRESCOTT, J. A. 1931. The soils of Australia in relation to vegetation and climate. Counc. Sci. Industr. Res. (Aust.) Bull. 52.

———————. 1934. Single value climatic factors. Trans. Proc. Roy. Soc. S.Aust. LVIII: 48-60.

———————. 1936. The climatic control of the Australian deserts. Trans. Proc. Roy. Soc. S.Aust. LX: 93-95.

———————. 1938. Indices in agricultural climatology. J. Aust. Inst. Agric. Sci. 4: 33-40.

———————. 1942. The phase and amplitude of the Australian mean monthly temperatures. Trans. Proc. Roy. Soc. S.Aust. LXVI: 46-49.

———————. 1943. A relationship between evaporation and temperature. Trans. Proc. Roy. Soc. S.Aust. LXVII: 1-6.

SZYMKIEWIEZ, D. 1925. In Acta Societatis Botanicorum Poloniae II: 239.

THORNTHWAITE, C. WARREN. 1931. The climates of North America according to a new classification. Geogr. Rev. 633-655.

———————. 1933. The climates of the earth. Geogr. Rev. 433-440.

TRUMBLE, H. C. 1939. Climatic factors in relation to the agricultural region of southern Australia. Trans. Proc. Roy. Soc. S.Aust. LXIII: 36-43.

ELECTROCARDIOGRAMS OF YOUNG LAMBS.

By W. T. KELSO, B.Sc., A.A.C.I., Analyst, Chemical Laboratory, Division of Plant Industry.

SUMMARY.

The heart rate of lambs shows a decrease with age. This decline is rapid for the first five weeks; between the fifth and ninth weeks the rate increases and from then to the twelfth week decreases. There is a tendency for all lambs to have the same proportional decrease by the end of the tenth week, even though the absolute heart rates are dissimilar.

The PR interval is independent of age, and minimum and maximum values show only small differences. Minimum and maximum values for QRS interval are more or less independent of age and show little variation. Variations in QT interval with age are more pronounced in mean, maximum, and minimum values. The value for this interval increases with age. PR and QT intervals vary inversely as the heart rate while QRS interval is fairly constant.

Indices derived from Bazett's systolic formula varied proportionately with heart rate and tended to a constant value with heart rates between 88 and 129 per minute.

Potentials of S, T, R, Q and QS were quite high, exceeding 500 microvolts in many instances. The P wave was much smaller. Instances of the absence of Q, R, S and QS waves were quite common.

Diphasic waves appeared in the T wave and to a lesser extent in P. The percentatge in the T wave was not great. P and R were invariably positive and Q, S and QS invariably negative. Serial changes in polarity of the T wave were mainly diphasic. Lead 1 (left foreleg to right foreleg) in the T wave was invariably positive. All waveforms were of predominant types with the usual exceptions. Transitional stages developed in serial recordings tending to the QS complex, occurring at approximately eight weeks. Lead 3 (left foreleg to left hindleg) was quite stable. Positive waves following the Q wave were not uncommon in Lead 3.

The ST function is discussed.

Average electrical axis changes about 10° every three weeks. Initial mean value for all lambs was −49°. Ten weeks later the mean was −127°. Changes in axis occurred in both directions with gradual progression towards negativity.

INTRODUCTION.

An instrument was designed to record responses from the administration of physiologically active substances to animals. Before any tests could be carried out it was necessary to collect relevant data from normal animals to decide what variations occurred with posture and age.

Merino sheep were selected as they represent the majority of animals subjected to investigation. Lambs were used because of economy and ease of handling. Their use also permitted the study of the effects of rumenal function on recordings at strategic periods.

Waveforms.

The electrical changes which precede auricular and ventricular contractions in the heart may be recorded as a series of waves which vary in frequency, shape, direction and potential. Anatomical variations, such as hypertrophy and position of the heart in the thoracic cavity, appear as corresponding variations in the nature of the waves.

This waveform, existing in its standard form (Plate 1, fig. 1) is divided into five sections designated P, Q, R, S and T. The base line represents the isoelectric line. Waves above and below this line are arbitrarily designated positive and negative, respectively.

The standard form is subject to variations and to avoid confusion the following nomenclature has been adopted throughout. Any downward deflection immediately following the P wave and followed by an upward deflection (however small) other than a T wave is designated as Q (Plate 1, fig. 2). Should an upward deflection (other than P) precede this downward deflection, the downward deflection is designated S (Plate 1, fig. 3). When no upward deflections (other than P or T) precede or succeed this downward deflection, the downward deflection is designated QS (Plate 1, fig. 4).

An appreciation of the differences between Q, S and QS explains the variations which occur in Tables 1-5. Positive and negative amplitudes are measured from the crest of each wave to the top and the bottom of the isoelectric line respectively. In general the PR interval is measured from the beginning of P to the beginning of Q; the QRS interval from the beginning of Q to the end of S; and the QT interval from the beginning of Q to the end of T.

In the study of electrocardiograms the time intervals and potentials of the main complexes, the shape of the waveform and any significant variations are the main concern.

APPARATUS.

The apparatus used in this work was designed by the author in association with T. A. Housley, V. W. Gibbs and I. Hamilton of the Brisbane staff of the Aeradio Division, Postmaster-General's Department, and constructed by the Division's staff.

A high-gain (voltage amplification of 2,000,000) low-frequency amplifier, with relatively flat response from zero frequency to 1 kilocycle, is used to amplify the potentials received. The frequency response and the time constants are such as to ensure faithful reproduction of the waveform which is being presented at the input. The amplified waveform is displayed

PREFACE TO PLATES.

The numerals (1, 2, 3) refer to the lead in question, while the letters (A, B, C, &c.) refer to the lamb used. Records adjoining one another denote the same lamb taken at a later period.



Plate 1.

Showing the standard waveform (Fig. 1) and its variations.



Plate 2.

Showing variations in waveform and amplitude with age, together with extremes of potential.

A, 5 weeks; B, 11 weeks; C, 3 weeks; D, 13 weeks; E, 2 weeks; F, 5 weeks; G, 1 week; H, 11 weeks; I, 1 week; J, 8 weeks; K, 4 weeks; L, 12 weeks; M, 3 weeks.



Plate 3.

Showing variations in polarity with age, together with extremes of potential.

A, 4 weeks; B, 15 weeks; C2, 8 weeks; C3, 5 weeks; D, 12 weeks; E, 1 week; F, 11 weeks; G, 2 weeks; H, 6 weeks; I, 5 weeks; J, 12 weeks; K, 2 weeks; L, 3 weeks; M, 5 weeks; N, 3 weeks; O, 10 weeks; P, 8 weeks.



Plate 4.

Showing variations in wave shape with age.

A1, 3, 5 weeks; A2, 2 weeks; B, 12 weeks; C, 3 weeks; D, 9 weeks; E, 2 weeks; F, 12 weeks; G, 2 weeks; H, 12 weeks; I, 5 weeks; J, 12 weeks; K, 5 weeks; L, 12 weeks.



Plate 5.

Showing variations in wave shape with age.

A, 4 weeks; B, 11 weeks; C, 1 week; D, 8 weeks; E, 2 weeks; F, 11 weeks; G, 2 weeks; H, 9 weeks; I, 7 weeks; J, 10 weeks; K, 1 week; L, 11 weeks.



Plate 6.

Showing the same lamb as in Plate 5 (D2, D3), lying on left side, standing, and lying on right side. All 8 weeks old.

on two cathode ray tubes having screen persistencies of the order of one minute in the dark. One cathode ray tube is used for photographic purposes only and the other for monitoring. No time base is used as the spot is photographed in the vertical direction only. The lens (f 4.5, focal length 2 inches) is arranged for a reduction of 2 : 1. Super XX film (35 mm.) moving uniformly at 25 mm. per second is used and the time lines shown thereon are produced by a neon lamp flashing 25 times a second and exposing the film through a slit 1/1,000 inch wide. A tachometer unit is incorporated to measure instantaneous and average heart rates up to 1,000 per minute.

EXPERIMENTAL.

Twelve Merino lambs, consisting of four males and eight females, were selected and recordings were commenced as soon as the lambs could with safety be separated from the mothers for the period necessary to complete all recordings. The number of ewes available for this experiment was fairly small, and since the lambs were born over a period of one month the number appearing in any one age group did not permit of statistical analysis of the data obtained.

Owing to the fact that very small alternating potentials from direct pickup are normally detrimental to the record, the animals were at first placed in an electrostatic shielded cage at earth potential. After a few recordings the animals outgrew the cage and subsequent results were obtained under unshielded conditions with the right hind leg earthed. Plates 2-6 show that the record suffered little distortion.

The animals were carried from their pens to the recording room and after about five minutes' rest the recordings were commenced. Since the procedure was carried out at about the same time each morning, the effect of handling, environmental temperature and humidity was reduced to a minimum. Throughout the recording period the animals were kept lying on the right side in order to prevent fibrillation of the isoelectric line arising from nervousness and muscle tremor. Plate 6 shows the same lamb recorded lying on the left side (M), standing (N) and lying on the right side (O). No marked variation, either in potential or in waveform, is seen in the recordings taken on the left and the right sides. In the standing position fibrillation of the isoelectric line obscures interpretation.

Standard electrode metal was used and contact was made with the skin by medium of sodium chloride jelly. The wool was clipped as short as possible and four electrodes were attached to the limbs, as shown below, to give the standard combinations:—

Lead 1.—Left foreleg to right foreleg.

Lead 2.—Left hindleg to right foreleg.

Lead 3.—Left hindleg to left foreleg.

Earth.—Right hindleg.

All recordings were examined by means of a projector using a magnification of 10; consequently, time intervals and potentials could be measured with a reasonable degree of accuracy. In all leads each determination was carried out 10 times and the mean taken. The vertical lines on the film are spaced 40 milliseconds apart and the calibrating potential appearing at the end of each record is usually 1 millivolt per 10 mm.

Calculation of the average electrical axis of the heart* has been attempted, and while the method employed (Dieuaide; see Ashman and Hull, 1944) is only approximate for absolute values, the comparison afforded some information.

Other methods which could be employed suffer a common disadvantage in that no correlation exists between anatomical structure, position and electrical considerations of the ruminant heart.

Bazett (1920) introduced the formula $K = \frac{QT}{\sqrt{C}}$ (where QT is the interval in seconds between beginning of Q to the end of T and C is the interval in seconds between consecutive R waves) to correlate the QT interval with the average heart rate cycle. Although primarily used to correct the QT interval in humans, in which case K is a constant, the formula is used to show the variation of K for a given QT interval and heart rate cycle.

DATA AND ANALYSIS.

Heart Rate.

Table 1 compares minimum, maximum and mean heart rates per minute with age. The table shows that there is a rapid decline in the mean rate from the second to the fifth week of life; however, between the fifth and ninth weeks the rate is higher than the level of the fifth week, but during the remaining three weeks of the test it is not markedly different from the rate of the fifth week.

The actual normal rates would, of course, be somewhat lower than the recorded rates, since handling would tend to increase the rate. The effect would be expected to decrease with the age of the animal.

No published information on the effect of temperature and humidity on young lambs is known, but Lee and Robinson (1941) in their work on mature sheep found that the heart rate tends to rise with high temperatures. It is also known that heart rate increases during digestion and with the level of the plane of nutrition and the rates for the fifth to the ninth week may possibly be explained as due to the fact that during this period the rumen is developing at the expense of the abomasum because of the larger intake of roughage.

---

* Determined from the variation in potentials of the QRS complex in Lead 1 and Lead 3. Variations occur mainly from ventricular hypertrophy and positions which the heart adopts in the thoracic cavity.

Table 1.

HEART RATES OF YOUNG LAMBS COMPARED WITH AGE.

Age to Nearest Week.	Number of Lambs.	Heart Rate per Minute. (Mean Value from 3 Leads.)		
		Min.	Max.	Mean.
1	2	183	217	200
2	7	189	231	208
3	3	147	179	161
4	2	141	157	149
5	6	115	160	136
6	1	150	150	150
7	3	131	170	144
8	5	152	192	167
9	3	141	194	163
10	2	114	140	127
11	2	133	141	137
12	5	88	140	122

Plate 5 (D2, D3) shows recordings taken in the morning for a lamb on its right side, after being deprived of food and water for 16 hours.

Plate 6, taken four hours later, shows recordings for the same lamb on its left (M2, M3) and right (O2, O3) sides and standing (N2, N3) after partaking of food and water. The heart rate in the morning is 140 and in the afternoon 115, while the T wave in Lead 2 has reversed its polarity and increased in potential.

Table 2.

HEART RATES OF YOUNG LAMBS SHOWING SERIAL VARIATIONS.

Lamb No.	Age at First Record in Weeks.	Heart Rate Per Min. at First Record.	Percentage Variation in Heart Rate* after—					
			3 Weeks.	4 Weeks.	6 Weeks.	7 Weeks.	8 Weeks.	10 Weeks.
507	1·5	216	−45	..	−22	..	..	−42
508	2·0	204	−32	..	− 6	..	..	−31
509	0·7	217	−35	..	−42	..	..	−35
510	1·4	222	−30	..	−31	..	..	−40
511	2·3	213	−25	..	−13	..	..	−37
512	2·6	179	−16	..	..	..	..	−36
513	2·0	194	..	..	..	−20	..	−37
514	1·7	189	−34	..	..	..	..	−53
521	1·5	231	..	−36	..	..	−34	..
522	3·0	160	..	+ 4	..	−40	..	..
523	1·4	183	..	−36	..	−23	..	..
524	3·0	147	..	−20	..	− 5	..	..
Mean Variation			−31	−22	−23	−22	−34	−39

* Variations are referred to the initial heart rate. Negative variations are decreases, positive variations are increases.

Table 2 shows the variation in heart rate occurring at specified intervals expressed as a percentage of the initial rate. It is seen from the table that, irrespective of age, the mean percentage variation after three weeks and 10 weeks is —31 and —39 respectively. During the period of age from five to nine weeks, five females and one male showed positive variations, one female a negative variation, and two males no variation, while data from three lambs (two females and one male) were not recorded.

Table 3.

SHOWING THE VARIATION OF PR, QRS, AND QT INTERVALS WITH RESPECT TO AGE.

Age to Nearest Week.	Number of Lambs.	PR			QRS			QT		
		Min.	Max.	Mean.	Min.	Max.	Mean.	Min.	Max.	Mean.
		secs.	secs.	secs.	secs.	secs.	secs.	secs.	secs.	secs.
1	1	·08	·08	·08	·04	·04	·04	·18	·18	·18
2	6	·03	·07	·06	·04	·06	·05	·17	·20	·19
3	3	·03	·07	·05	·04	·07	·05	·20	·22	·21
4	2	·06	·07	·06	·03	·05	·04	·22	·24	·23
5	6	·06	·10	·08	·03	·05	·04	·19	·26	·24
6	1	·08	·08	·08	·02	·04	·03	·12	·20	·17
7	4	·07	·08	·08	·04	·04	·04	·19	·24	·22
8	5	·06	·09	·07	·02	·05	·03	·18	·23	·20
9	2	·08	·08	·08	·03	·03	·03	·23	·23	·23
10	2	·08	·08	·08	·04	·04	·04	·22	·24	·23
11	2	·08	·08	·08	·04	·04	·04	·21	·22	·22
12	5	·08	·12	·09	·04	·04	·04	·22	·28	·25

Table 4.

SHOWING THE VARIATION OF PR, QRS, QT, AND K WITH HEART RATE.

Heart Rate Range.	Number of Lambs.	PR			QT			QRS			Number of Lambs.	K		
		Min.	Max.	Mean.	Min.	Max.	Mean.	Min.	Max.	Mean.		Min.	Max.	Mean.
240–200	18	·04	·07	·06	·15	·21	·18	·04	·08	·05	3	·350	·380	·367
199–170	20	·02	·08	·07	·16	·25	·19	·02	·09	·04	7	·305	·442	·358
169–130	55	·06	·09	·07	·18	·25	·21	·02	·06	·04	20	·288	·383	·336
129– 88	27	·07	·11	·08	·21	·27	·24	·02	·07	·04	10	·326	·379	·341

Intervals.

All intervals are measured in accordance with the procedure given on page 68. Table 3 sets out minimum, maximum and mean values of the PR, QRS, and QT intervals with respect to age. In all cases any one value is the mean of three leads.

Table 4 shows how PR, QRS, QT and K vary with heart rate.

1. PR Interval.

The PR interval remains fairly constant with age. Table 3 shows very little difference between minimum and maximum values. However, the interval was not quite the same for all leads, being greatest in Lead 3 with Leads 1 and 2 about equal. These differences were of the order of 0.01 sec. and instances of Lead 1 being greater than Lead 3 were extremely rare. In very few instances was the Q wave apparent, and, associated with the inter-relationship which exists between Q, S and QS, the PR interval (measured from beginning of P to the beginning of the next deflection whether +ve or —ve) was found to be fairly constant as shown from the minimum values for any one age group.

Table 4 shows that the mean value of the PR interval exhibits a regular increase in duration with decreasing heart rates. The same variations apply to the maximum values and to a lesser extent to the minimum values.

2. QRS Interval.

The relative constancy of the duration of the QRS interval should be noted (Table 3). In several instances no difference existed between the minimum and the maximum values. The constancy of this duration exists in spite of the variation in waveform which was manifest from one serial record to another. When transitional waveforms (i.e., interchanging of Q, S and QS) appeared, this interval was measured from the beginning of Q, R and QS to the end of R, S, and QS respectively. When Q waves were in evidence, they were relatively large. In only a few instances did small Q waves exist, and consequently quite a large number of leads showed no Q waves at all. When this occurred the time interval was measured from the first wave following the P wave. As a general rule, the value of the duration was greatest in Lead 1, while Leads 2 and 3 were almost equal.

Table 4 shows that the mean duration of the QRS interval is independent of the heart rate. The minima and the maxima are widely divergent.

3. QT Interval.

A progressive increase with age is noted in the QT interval with the exception of anomalies occurring between the sixth and ninth weeks (Table 3). The minimum and maximum values as a whole show relatively small differences. In general the greatest value was obtained in Lead 1, with Lead 2 greater than Lead 3. Even then the differences between Leads 1, 2 and 3 were proportionately small. The variation between leads was more definite than in the PR and QRS intervals. In the QT interval, as with the QRS interval above, the same problem of transitional Q, S and QS waves is encountered together with the absence of Q waves. Coupled with this measurement there are depressed and elevated ST junctions (Plates 2(A1) and 5(C2)) and also diphasic T waves (Plate 3(O3)). ST junction variations and diphasic T waves


FREQUENCY DISTRIBUTION OF POTENTIAL
P
Q
R
%
1
2
3



Figure 1.

Showing frequency distributions of potential. All potentials in microvolts. Number of determinations, 40.

amounted to 25 per cent. and 13 per cent. of the number of records respectively. Serial variations of the T wave from monophasic to diphasic forms were not a predominant feature.

In Table 4, minimum, maximum and mean values are shown to vary inversely with the heart rate in a regular manner. The same result is noticed in humans (Pardee, 1943). The regular variations in this group are more definite than in either of the two preceding groups.

4. K (Systolic Index).

The small potentials which appeared regularly in Lead 3 rendered it unsuitable for calculation of K. In Lead 1 the QRS complex was more stable and subject to less serial variations than in Lead 2, and consequently determinations were made in this lead.

The mean value of K for various heart rate ranges is shown in Table 4. Over the wide heart rate range, the change in the value of K is not large, and there is a tendency to maintain a constant value. It is not possible to state accurately the manner in which K varies with heart rate until the relationship between the QT interval and heart rate is known. Alfredson and Sykes (1942) found K to vary inversely with heart rate. A more detailed mathematical study is necessary to establish the true relationship.

Table 5.

Showing the Potentials in all Leads.

Deflection.	Lead.	Potential in Microvolts.			Deflection.	Lead.	Potential in Microvolts.		
		Min.	Max.	Mean.			Min.	Max.	Mean.
P	1	0	267	118	S	1	0	893	427
	2	60	235	144		2	0	1,400	360
	3	0	180	72		3	0	1,187	88
Q	1	0	135	17	T	1	38	620	202
	2	0	910	107		2	18	540	177
	3	77	1,000	183		3	33	408	112
R	1	0	941	200	QS	1	0	887	120
	2	0	750	163		2	0	1,236	166
	3	Trace	936	233		3	0	996	91

Potentials.

Table 5 shows the variation in potential of P, Q, R, S, T and QS for all leads. The figures show no marked tendency to conform to any general form and wide variations in potential occurred just as in humans. The results are expressed in the tables in microvolts purely for convenience. Effect of

position of the animal when the record was taken is not considerable (see Plate 6, M1, M2, M3; O1, O2, O3) and skin contacts should be reasonably constant since standard technique was used throughout. Nil results were prevalent in Q, R, S and QS in nearly all leads, while in P and T deflections measurable values predominated in all leads. Low values are present in Lead 3 in P, S, T and QS deflections. Q, R, S and QS are conspicuous for their large maximum values.

Figure 1 shows the frequency distribution of potential in any one lead. For all leads in the Q, S and QS group a large proportion of nil recordings was obtained. This was brought about in the main by transitional waveforms. P and T waves are singularly free from nil results in all leads.

The potentials of the S and P waves were respectively the largest and smallest recorded. Typical illustrations of the following waveforms are presented:—Large P—Plate 2 (M2); P absent—Plate 2 (M3); large Q—Plate 2 (A1) and Plate 3 (A3); large T—Plate 3 (I1); and large R and S—Plate 5 (C2).

Wave Form Classification.

P Wave.

The P wave is predominantly positive, but a few diphasic waves were encountered (Plate 2 (B1, B3)). The predominant feature is the appearance of this slightly pointed waveform in all leads in 70 per cent. of the recordings. Some waves are relatively small and these occur mainly in Lead 3. Minor variations such as "steep front" and "slurred back" and "sharp front" and "sharp back" amount to about 12 per cent. Illustrations of these various wave forms appear in Plate 2. The variations when present are spread throughout all leads and occur at all ages. Serial records reveal slight changes.

QRS Complex.

From the definition of R, Q and S, 50 per cent. of the waves fall in the very small and nil category. This is the result of the inter-relationship existing within this group. The general wave shape took the form of pointed equal sloped waves, which amounted to about 25 per cent. Leads 1 and 2 were characterized by their number of nil waves. Lead 3 was not predominant in this respect. Positive waves immediately preceding a T wave amounted to 15 per cent. of the total number of recordings. Illustrations of the various waveforms encountered are seen in Plates 4 and 5. In general, the downward stroke is much thicker than the forward stroke. Lead 3 is well represented in the various waveform groupings.

The change in the QRS group with serial recordings is noteworthy. The initial waveform for each animal conforms to the standard P, Q, R, S and T pattern for humans except that the Q wave is not well pronounced and is more often zero. The R wave reduces in amplitude to zero and the original S wave now becomes QS (see Plates 4 (A1, B1; A2, B2; E1, F1; C1, D1). In Lead 1 this QS complex occurs eight times at an average age of eight weeks.

It occurs six times in Lead 2 with an average age of six weeks. QS did not appear in Lead 3 where the predominant feature was the appearance of a +ve wave immediately preceding the T wave. Lead 3 can be regarded as the most stable so far as this experiment is concerned. The QS wave shown in Plate 4 (G2) occurred initially and tended to remain constant throughout. Plate 5 (G1 and H1) and Plate 4 (K2 and L2) show still another variation. In only one instance did a QS wave occurring initially change to another form (see Plate 5 (I2 and J2)).

Miscellaneous variations of the QRS group are illustrated in Plates 4 and 5. The tendency in these changes is towards the QS complex, which appears to be the stable form in all leads. The illustrations are as follows:— Lead 1—Plate 4 (IJ and GH); Lead 2—Plate 4 (EF and CD); and Plate 5 (CD and AB); Lead 3—Plate 4 (AB, EF, CD and GH) and Plate 5 (CD, EF, KL and AB).

T Wave and ST Junction.

Three types of polarity exist, viz., positive, negative and diphasic. Positive waves occur in 64 per cent. of the leads, negative waves in 22 per cent. and diphasic waves in 14 per cent. Lead 1 is composed wholly of positive waves, while in Leads 2 and 3 positive and negative waves respectively predominate. When diphasic waves occur they are always of the negative-positive type. Twenty-three variations, from one form to another, occurred in the serial waveforms. About one-half of these variations were of diphasic type, and only on three occasions did a diphasic wave change to any other form. Two lambs in the initial record had diphasic waves. T waves are characterized by a slurred front and steeper back. Some waves have rounded peaks but the majority are pointed.

Plate 3 shows variations in polarity with age together with extremes of potential. Large potentials are seen in P1 and P2, and small values in L1, M2 and N3. Pointed and slurred waves are illustrated in I1 and J1, respectively. A general feature of the T wave is that its form is determined by the polarity of the preceding wave. If this is positive then the polarity of T is mainly diphasic, with instances of a few negative values. If the preceding wave is negative then the T wave is invariably +ve. These instances have been illustrated in Plate 3.

The ST junction is the mode in which the T wave leaves the isoelectric line. About 75 per cent. of the junctions commence at the isoelectric line, 14 per cent. below and 10 per cent. above. For heart rates of about 200 per minute, the T wave arises from the S wave, there being no isoelectric period (Plate 3 (P1)). Changes in ST junctions occurred 23 times, of which three were double changes. No general principle was involved in the sequence of these changes.

Average Electrical Axis.

Values calculated from the initial record range from −60° to +15°, corresponding to a range of 75° measured in the direction of decreasing negative and positive values. The mean value is −49°. Ten weeks later the minimum

and maximuum values were −40° and −168° respectively, with a mean of −127°. With increasing age the general trend is for the axis value to change to a more negative value. So far as the individual lambs were concerned, the serial variations were quite interesting. The change amounted to approximately 10° every three weeks.

All changes did not occur in the same direction and in several instances the variations in any one lamb were such that the ultimate direction was that of decreased negativity. It was possible to calculate the axis in practically all cases.

From Plate 6 it can be seen that position has very little effect on the axis at the age of eight weeks. More marked deviations could be expected in the earlier stages of life.

CONCLUSIONS.

The data suggest that it is unwise to select lambs under the age of three months for the purpose of demonstrating induced electrocardiographic changes. Lambs three months of age and older are relatively stable in behaviour and in addition they lend themselves to experimental procedure with the minimum amount of handling. Recordings can be taken without the aid of a screened room in all positions other than standing, but the use of the room is strongly advisable.

ACKNOWLEDGEMENTS.

The author is indebted to Dr. M. White, Agricultural Chemist, who initiated the project, arranged for the production of the equipment and gave helpful criticism; to Prof. D. H. K. Lee, Professor of Physiology, University of Queensland, for providing facilities; to Dr. F. H. S. Roberts, formerly Acting Director of Research in the Department's Division of Animal Industry, for supplying the experimental animals; to Messrs. R. Riek, G. Klemm, and A. Tuffley of the University of Queensland Department of Physiology, and to Mr. W. Manley, Illustrator in the Division of Plant Industry, for valuable assistance.

REFERENCES

ALFREDSON, B. V., and SYKES, J. F. 1942. Electrocardiograph studies in normal dairy cattle. J.Agric.Res. 65: 61-87.

ASHMAN, R., and HULL, E. 1944. Essentials of Cardiography. New York: MacMillan and Co.

BAZETT, H. C. 1920. An analysis of the time relations in electrocardiograms. Heart 7: 353-370.

GOLDBERG, H. 1944. Bio-electric research apparatus. Proc. Inst. Ref. Eng. 32: 330.

LEE, D. H. K., and ROBINSON, K. 1941. Reactions of the sheep to hot atmospheres. Proc. Roy. Soc. Qld. LIII: 189-200.

PARDEE, H. E. B. 1943. Clinical Aspects of the Electrocardiogram. New York: Hoeber.

PARR and WALTER. 1943. Amplifying and recording technique in electro-biology with species reference to the electrical activity of the human brain. J.Inst.Elec.Eng. 90: 129.

STURM, R. E., and WOOD, E. H. An instantaneously recording cardiotachometer. Personal communication.

A. H. TUCKER, Government Printer, Brisbane.